

## CRP U-hs \*

Diagnostic reagent for quantitative in vitro determination of C-reactive Protein (CRP) in serum or plasma on DiaSys respons<sup>®</sup>920

### Order information

Cat. No. 1 7045 99 10 920

4 twin containers for 100 determinations each

### Method

Particle enhanced immunoturbidimetric test with two applications. The high sensitive application is recommended for samples with concentrations lower than 20 mg/L and where high precision and extremely good sensitivity is required (measuring range of hs application: 0.05 mg/L – 20 mg/L). The universal application is characterized by a wide measuring range (2.5 – 350 mg/L) and low sample volume.

### Principle

Determination of the CRP concentration by photometric measurement of antigen-antibody reaction of antibodies to human CRP bound to polystyrene particles with CRP present in the sample.

### Reagents

#### Components and Concentrations

<b>R1:</b>	HEPES	pH 7.2	10 mmol/L
<b>R2:</b>	Borate buffer		4.6 mmol/L
	Polyclonal (goat) and monoclonal (mouse) anti-human CRP antibodies bound to carboxylated polystyrene particles		

#### Storage Instructions and Reagent Stability

The reagents are stable up to the end of the indicated month of expiry, if stored at 2 – 8°C, protected from light and contamination is avoided. DiaSys respons containers provide protection from light. Do not freeze the reagents!

#### Warnings and Precautions

- The reagents contain sodium azide (0.95 g/L) as preservative. Do not swallow! Avoid contact with skin and mucous membranes.
- To avoid carryover interference, please take care of efficient washing especially after use of interfering reagents. Please refer to the DiaSys respons<sup>®</sup>920 Carryover Pair Table. Carryover pairs and automated washing steps with the recommended cleaning solution can be specified in the system software. Please refer to the user manual.
- In very rare cases, samples of patients with gammopathy might give falsified results [9].
- Heterophile antibodies in patient samples may cause falsified results.
- Please refer to the safety data sheets and take the necessary precautions for the use of laboratory reagents. For diagnostic purposes, the results should always be assessed with the patient's medical history, clinical examinations and other findings.
- For professional use only!

#### Waste Management

Please refer to local legal requirements.

#### Reagent Preparation

The reagents are ready to use. Do not shake and avoid foaming. The bottles are placed directly into the reagent rotor.

### Specimen

Serum, heparin plasma or EDTA plasma

Stability [1]:	15 days	at	20 – 25°C
	2 months	at	4 – 8°C
	3 years	at	-20°C

Only freeze once! Discard contaminated specimens.

### Calibrators and Controls

DiaSys TruCal CRP U calibrator set (universal range) or DiaSys TruCal CRP hs calibrator set (high-sensitive range) are recommended for calibration. The assigned values of the calibrators have been made traceable to the IFCC reference material ERM<sup>®</sup>-DA474/IFCC. For internal quality control, DiaSys TruLab CRP or TruLab Protein (universal range) or TruLab CRP hs (high sensitive range) controls should be assayed. Each laboratory should establish corrective action in case of deviations in control recovery.

	Cat. No.	Kit size
TruCal CRP U (5 Level)	1 7040 99 10 059	5 x 1 mL
TruCal CRP hs (5 Level)	1 7080 99 10 059	5 x 1 mL
TruLab CRP hs Level 1	5 9730 99 10 046	3 x 1 mL
TruLab CRP hs Level 2	5 9740 99 10 046	3 x 1 mL
TruLab CRP Level 1	5 9600 99 10 045	3 x 2 mL
TruLab CRP Level 2	5 9610 99 10 045	3 x 2 mL
TruLab Protein Level 1	5 9500 99 10 046	3 x 1 mL
TruLab Protein Level 2	5 9510 99 10 046	3 x 1 mL

### Performance Characteristics Universal Application

Measuring range from 2.5 to 350 mg/L CRP, at least up to the concentration of the highest calibrator (in case of higher concentrations re-measure samples after manual dilution with NaCl solution (9 g/L) or use the rerun function).

Limit of detection**	0.8 mg/L CRP
No prozone effect up to 1000 mg/L CRP	
On-board stability	8 weeks
Calibration stability	6 weeks

Interfering substance	Interferences < 10%	CRP [mg/L]
<b>Hemoglobin</b>	up to 700 mg/dL	4.15
	up to 1200 mg/dL	40.7
<b>Bilirubin, conjugated</b>	up to 70 mg/dL	4.15
	up to 70 mg/dL	41.6
<b>Bilirubin, unconjugated</b>	up to 60 mg/dL	4.21
	up to 60 mg/dL	42.6
<b>Lipemia (triglycerides)</b>	up to 2100 mg/dL	3.96
	up to 2100 mg/dL	41.0
<b>Rheumatoidfactor</b>	up to 600 IU/mL	4.41

For further information on interfering substances refer to Young DS [8].

Precision			
Within run (n=20)	Sample 1	Sample 2	Sample 3
Mean [mg/L]	5.68	11.7	68.0
Coefficient of variation [%]	2.13	1.51	1.99
Between run (n=20)	Sample 1	Sample 2	Sample 3
Mean [mg/L]	4.74	11.8	68.6
Coefficient of variation [%]	3.72	2.30	2.91

Method comparison (n=153)	
Test x	DiaSys CRP U-hs (Hitachi 917)
Test y	DiaSys CRP U-hs (respons <sup>®</sup> 920)
Slope	1.01
Intercept	0.123 mg/L
Coefficient of correlation	0.994

\*\* according to NCCLS document EP17-A, vol. 24, no. 34

### Performance Characteristics high sensitive Application

Measuring range from 0.05 to 20 mg/L CRP, at least up to the concentration of the highest calibrator (in case of higher concentrations re-measure samples after manual dilution with NaCl solution (9 g/L) or use the rerun function).

Limit of detection**	0.05 mg/L CRP
No prozone effect up to 800 mg/L CRP	
On-board stability	7 weeks
Calibration stability	3 weeks

Interfering substance	Interferences < 10%	CRP [mg/L]
<b>Lipemia (triglycerides)</b>	up to 1450 mg/dL	1.29

For further information on interfering substances refer to Young DS [8].

Precision			
Within run (n=20)	Sample 1	Sample 2	Sample 3
Mean [mg/L]	0.38	2.20	18.0
Coefficient of variation [%]	3.35	0.97	1.44
Between run (n=20)	Sample 1	Sample 2	Sample 3
Mean [mg/L]	0.74	1.80	11.5
Coefficient of variation [%]	2.46	1.68	2.80

Method comparison (n=111)	
Test x	DiaSys CRP U-hs (Hitachi 911)
Test y	DiaSys CRP U-hs (respons <sup>®</sup> 920)
Slope	1.02
Intercept	0.042 mg/L
Coefficient of correlation	0.999

\*\* lowest measurable activity which can be distinguished from zero mean + 3 SD (n=20) of an analyte free specimen

## Reference Range [2,3]

Adults	< 5 mg/L
Newborns up to 3 weeks	< 4.1 mg/L
Infants and children	< 2.8 mg/L

Each laboratory should check if the reference ranges are transferable to its own patient population and determine own reference ranges if necessary.

## Literature

1. Guder WG, Zawta B et al. The Quality of Diagnostic Samples. 1<sup>st</sup> ed. Darmstadt: GIT Verlag; 2001; p. 24-5.
2. Dati F, Schumann G, Thomas L, Aguzzi F, Baudner S, Bienvenu J et al. Consensus of a group of professional societies and diagnostic companies on guidelines for interim reference ranges for 14 proteins in serum based on the standardization against the IFCC/BCR/CAP reference material (CRM 470). Eur J Clin Chem Clin Biochem 1996; 34: 517-20.
3. Schiebusch H, Liappis N, Klein G. High sensitive CRP and creatinine: reference intervals from infancy to childhood. Poster präsentiert am AACC/CSCC; Juli/August 2001, Chicago, Illinois.
4. Thompson D, Milford-Ward A, Whicher JT. The value of acute phase protein measurements in clinical practice. Ann Clin Biochem 1992; 29: 123-31.
5. Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. N Engl J Med 1999; 340: 448-54.
6. Hansson LO, Lindquist L. C-reactive protein: its role in the diagnosis and follow-up of infectious diseases. Curr Opin Infect Diseases 1997; 10: 196-201.
7. Sipe JD. Acute-phase proteins in osteoarthritis. Semin Arthritis Rheum 1995; 25: 75-86.
8. Young DS. Effects of Drugs on Clinical Laboratory Tests. 5<sup>th</sup> ed. Volume 1 and 2. Washington, DC: The American Association for Clinical Chemistry Press 2000.
9. Bakker AJ, Mücke M. Gammopathy interference in clinical chemistry assays: mechanisms, detection and prevention. ClinChemLabMed 2007;45(9):1240-1243.

## Hersteller



DiaSys Diagnostic Systems GmbH  
Alte Strasse 9 65558 Holzheim Deutschland

## CRP U-hs

Universal application for serum und plasma

Test Details		Test Volumes		Reference Ranges	
Test	: CRPU			Auto Rerun	<input type="checkbox"/>
Report Name	: C-Reactive Protein Universal			Online Calibration	<input type="checkbox"/>
Unit	: mg/L	Decimal Places	: 2	Cuvette Wash	<input type="checkbox"/>
Wavelength-Primary	: 505	Secondary	: 0	Total Reagents	: 2
Assay Type	: 2-Point	Curve Type	: 4P Logit Log	Reagent R1	: CRP U-hs R1
M1 Start	: 19	M1 End	: 19	Reagent R2	: CRP U-hs R2
M2 Start	: 31	M2 End	: 31	<b>Consumables/Calibrators:</b>	
Sample Replicates	: 1	Standard Replicates	: 3	Blank /Level 0	: 0
Control Replicates	: 1	Control Interval	: 0	Calibrator 1	: **
Reaction Direction	: Increasing	React. Abs. Limit	: *	Calibrator 2	: **
Prozone Limit %	: 97	Prozone Check	: Lower	Calibrator 3	: **
Linearity Limit %	: 0	Delta Abs. / Min.	: 0.00	Calibrator 4	: **
Technical Minimum	: *	Technical Maximum	: *	Calibrator 5	: **
Y = aX + b	a= 1.00	b=	: 0.00		

\*Technical limits are automatically defined by the software by the upper and lower calibrator level.

\*\* Enter calibrator value.

Test Details		Test Volumes		Reference Ranges																																			
Test	: CRPU																																						
Sample Type	: Serum																																						
<table border="1"> <thead> <tr> <th colspan="4">Sample Volumes</th> </tr> </thead> <tbody> <tr> <td>Normal</td> <td>: 2.0 µL</td> <td>Dilution Ratio</td> <td>: 1 X</td> </tr> <tr> <td>Increase</td> <td>: 4.0 µL</td> <td>Dilution Ratio</td> <td>: 1 X</td> </tr> <tr> <td>Decrease</td> <td>: 2.0 µL</td> <td>Dilution Ratio</td> <td>: 6 X</td> </tr> <tr> <td>Standard Volume</td> <td>: 2.0 µL</td> <td></td> <td></td> </tr> </tbody> </table>				Sample Volumes				Normal	: 2.0 µL	Dilution Ratio	: 1 X	Increase	: 4.0 µL	Dilution Ratio	: 1 X	Decrease	: 2.0 µL	Dilution Ratio	: 6 X	Standard Volume	: 2.0 µL			<table border="1"> <thead> <tr> <th colspan="2">Sample Types</th> </tr> </thead> <tbody> <tr> <td><input checked="" type="checkbox"/></td> <td>Serum</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Urine</td> </tr> <tr> <td><input type="checkbox"/></td> <td>CSF</td> </tr> <tr> <td><input checked="" type="checkbox"/></td> <td>Plasma</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Whole Blood</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Other</td> </tr> </tbody> </table>		Sample Types		<input checked="" type="checkbox"/>	Serum	<input type="checkbox"/>	Urine	<input type="checkbox"/>	CSF	<input checked="" type="checkbox"/>	Plasma	<input type="checkbox"/>	Whole Blood	<input type="checkbox"/>	Other
Sample Volumes																																							
Normal	: 2.0 µL	Dilution Ratio	: 1 X																																				
Increase	: 4.0 µL	Dilution Ratio	: 1 X																																				
Decrease	: 2.0 µL	Dilution Ratio	: 6 X																																				
Standard Volume	: 2.0 µL																																						
Sample Types																																							
<input checked="" type="checkbox"/>	Serum																																						
<input type="checkbox"/>	Urine																																						
<input type="checkbox"/>	CSF																																						
<input checked="" type="checkbox"/>	Plasma																																						
<input type="checkbox"/>	Whole Blood																																						
<input type="checkbox"/>	Other																																						
<table border="1"> <thead> <tr> <th colspan="4">Reagent Volumes and Stirrer Speed</th> </tr> </thead> <tbody> <tr> <td>RGT-1 Volume</td> <td>: 120 µL</td> <td>R1 Stirrer Speed</td> <td>: Medium</td> </tr> <tr> <td>RGT-2 Volume</td> <td>: 120 µL</td> <td>R2 Stirrer Speed</td> <td>: Medium</td> </tr> </tbody> </table>				Reagent Volumes and Stirrer Speed				RGT-1 Volume	: 120 µL	R1 Stirrer Speed	: Medium	RGT-2 Volume	: 120 µL	R2 Stirrer Speed	: Medium																								
Reagent Volumes and Stirrer Speed																																							
RGT-1 Volume	: 120 µL	R1 Stirrer Speed	: Medium																																				
RGT-2 Volume	: 120 µL	R2 Stirrer Speed	: Medium																																				

Test Details		Test Volumes		Reference Ranges																																			
Test	: CRPU																																						
Sample Type	: Serum																																						
Reference Range	: DEFAULT																																						
Category	: Male																																						
<table border="1"> <thead> <tr> <th colspan="4">Reference Range</th> </tr> <tr> <td></td> <td>Lower Limit</td> <td>Upper Limit</td> <td></td> </tr> <tr> <td></td> <td>(mg/L)</td> <td>(mg/L)</td> <td></td> </tr> </thead> <tbody> <tr> <td>Normal</td> <td>: 0.00</td> <td>: 5.00</td> <td></td> </tr> <tr> <td>Panic</td> <td>: 0.00</td> <td>: 0.00</td> <td></td> </tr> </tbody> </table>				Reference Range					Lower Limit	Upper Limit			(mg/L)	(mg/L)		Normal	: 0.00	: 5.00		Panic	: 0.00	: 0.00		<table border="1"> <thead> <tr> <th colspan="2">Sample Types</th> </tr> </thead> <tbody> <tr> <td><input checked="" type="checkbox"/></td> <td>Serum</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Urine</td> </tr> <tr> <td><input type="checkbox"/></td> <td>CSF</td> </tr> <tr> <td><input checked="" type="checkbox"/></td> <td>Plasma</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Whole Blood</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Other</td> </tr> </tbody> </table>		Sample Types		<input checked="" type="checkbox"/>	Serum	<input type="checkbox"/>	Urine	<input type="checkbox"/>	CSF	<input checked="" type="checkbox"/>	Plasma	<input type="checkbox"/>	Whole Blood	<input type="checkbox"/>	Other
Reference Range																																							
	Lower Limit	Upper Limit																																					
	(mg/L)	(mg/L)																																					
Normal	: 0.00	: 5.00																																					
Panic	: 0.00	: 0.00																																					
Sample Types																																							
<input checked="" type="checkbox"/>	Serum																																						
<input type="checkbox"/>	Urine																																						
<input type="checkbox"/>	CSF																																						
<input checked="" type="checkbox"/>	Plasma																																						
<input type="checkbox"/>	Whole Blood																																						
<input type="checkbox"/>	Other																																						

## CRP U-hs

### hs application for Serum and Plasma

Test Details	Test Volumes	Reference Ranges
Test : CRPhs		Auto Rerun : <input type="checkbox"/>
Report Name : CRP high sensitive		Online Calibration <input type="checkbox"/>
Unit : mg/L	Decimal Places : 3	Cuvette Wash <input type="checkbox"/>
Wavelength-Primary : 505	Secondary : 0	Total Reagents : 2
Assay Type : 2-Point	Curve Type : Cubic Spline	Reagent R1 : CRP U-hs R1
M1 Start : 19	M1 End : 19	Reagent R2 : CRP U-hs R2
M2 Start : 33	M2 End : 33	<b>Consumables/Calibrators:</b>
Sample Replicates : 1	Standard Replicates : 3	Blank /Level 0 : 0
Control Replicates : 1	Control Interval : 0	Calibrator 1 : **
Reaction Direction : Increasing	React. Abs. Limit : *	Calibrator 2 : **
Prozone Limit % : 97	Prozone Check : Lower	Calibrator 3 : **
Linearity Limit % : 0	Delta Abs. / Min. : 0.00	Calibrator 4 : **
Technical Minimum : *	Technical Maximum : *	Calibrator 5 : **
Y = aX + b    a= : 1.00	b= : 0.00	

\*Technical limits are automatically defined by the software via the upper and lower calibrator level.

\*\* Enter calibrator value.

Test Details	Test Volumes	Reference Ranges																											
Test : CRPhs																													
Sample Type : Serum																													
<table border="1"> <thead> <tr> <th colspan="4">Sample Volumes</th> </tr> </thead> <tbody> <tr> <td>Normal</td> <td>: 10.00 <math>\mu</math>L</td> <td>Dilution Ratio</td> <td>: 1 X</td> </tr> <tr> <td>Increase</td> <td>: 20.00 <math>\mu</math>L</td> <td>Dilution Ratio</td> <td>: 1 X</td> </tr> <tr> <td>Decrease</td> <td>: 5.00 <math>\mu</math>L</td> <td>Dilution Ratio</td> <td>: 1 X</td> </tr> <tr> <td>Standard Volume</td> <td>: 10.00 <math>\mu</math>L</td> <td></td> <td></td> </tr> </tbody> </table>		Sample Volumes				Normal	: 10.00 $\mu$ L	Dilution Ratio	: 1 X	Increase	: 20.00 $\mu$ L	Dilution Ratio	: 1 X	Decrease	: 5.00 $\mu$ L	Dilution Ratio	: 1 X	Standard Volume	: 10.00 $\mu$ L			<table border="1"> <thead> <tr> <th>Sample Types</th> </tr> </thead> <tbody> <tr> <td><input checked="" type="checkbox"/> Serum</td> </tr> <tr> <td><input type="checkbox"/> Urine</td> </tr> <tr> <td><input type="checkbox"/> CSF</td> </tr> <tr> <td><input checked="" type="checkbox"/> Plasma</td> </tr> <tr> <td><input type="checkbox"/> Whole Blood</td> </tr> <tr> <td><input type="checkbox"/> Other</td> </tr> </tbody> </table>	Sample Types	<input checked="" type="checkbox"/> Serum	<input type="checkbox"/> Urine	<input type="checkbox"/> CSF	<input checked="" type="checkbox"/> Plasma	<input type="checkbox"/> Whole Blood	<input type="checkbox"/> Other
Sample Volumes																													
Normal	: 10.00 $\mu$ L	Dilution Ratio	: 1 X																										
Increase	: 20.00 $\mu$ L	Dilution Ratio	: 1 X																										
Decrease	: 5.00 $\mu$ L	Dilution Ratio	: 1 X																										
Standard Volume	: 10.00 $\mu$ L																												
Sample Types																													
<input checked="" type="checkbox"/> Serum																													
<input type="checkbox"/> Urine																													
<input type="checkbox"/> CSF																													
<input checked="" type="checkbox"/> Plasma																													
<input type="checkbox"/> Whole Blood																													
<input type="checkbox"/> Other																													
<table border="1"> <thead> <tr> <th colspan="4">Reagent Volumes and Stirrer Speed</th> </tr> </thead> <tbody> <tr> <td>RGT-1 Volume</td> <td>: 90 <math>\mu</math>L</td> <td>R1 Stirrer Speed</td> <td>: Medium</td> </tr> <tr> <td>RGT-2 Volume</td> <td>: 90 <math>\mu</math>L</td> <td>R2 Stirrer Speed</td> <td>: Medium</td> </tr> </tbody> </table>		Reagent Volumes and Stirrer Speed				RGT-1 Volume	: 90 $\mu$ L	R1 Stirrer Speed	: Medium	RGT-2 Volume	: 90 $\mu$ L	R2 Stirrer Speed	: Medium																
Reagent Volumes and Stirrer Speed																													
RGT-1 Volume	: 90 $\mu$ L	R1 Stirrer Speed	: Medium																										
RGT-2 Volume	: 90 $\mu$ L	R2 Stirrer Speed	: Medium																										

Test Details	Test Volumes	Reference Ranges																											
Test : CRPhs																													
Sample Type : Serum																													
Reference Range : DEFAULT																													
Category : Male																													
<table border="1"> <thead> <tr> <th colspan="4">Reference Range</th> </tr> <tr> <th></th> <th>Lower Limit</th> <th>Upper Limit</th> <th></th> </tr> <tr> <th></th> <th>(mg/L)</th> <th>(mg/L)</th> <th></th> </tr> </thead> <tbody> <tr> <td>Normal</td> <td>: 0.00</td> <td>: 5.00</td> <td></td> </tr> <tr> <td>Panic</td> <td>: 0.00</td> <td>: 0.00</td> <td></td> </tr> </tbody> </table>		Reference Range					Lower Limit	Upper Limit			(mg/L)	(mg/L)		Normal	: 0.00	: 5.00		Panic	: 0.00	: 0.00		<table border="1"> <thead> <tr> <th>Sample Types</th> </tr> </thead> <tbody> <tr> <td><input checked="" type="checkbox"/> Serum</td> </tr> <tr> <td><input type="checkbox"/> Urine</td> </tr> <tr> <td><input type="checkbox"/> CSF</td> </tr> <tr> <td><input checked="" type="checkbox"/> Plasma</td> </tr> <tr> <td><input type="checkbox"/> Whole Blood</td> </tr> <tr> <td><input type="checkbox"/> Other</td> </tr> </tbody> </table>	Sample Types	<input checked="" type="checkbox"/> Serum	<input type="checkbox"/> Urine	<input type="checkbox"/> CSF	<input checked="" type="checkbox"/> Plasma	<input type="checkbox"/> Whole Blood	<input type="checkbox"/> Other
Reference Range																													
	Lower Limit	Upper Limit																											
	(mg/L)	(mg/L)																											
Normal	: 0.00	: 5.00																											
Panic	: 0.00	: 0.00																											
Sample Types																													
<input checked="" type="checkbox"/> Serum																													
<input type="checkbox"/> Urine																													
<input type="checkbox"/> CSF																													
<input checked="" type="checkbox"/> Plasma																													
<input type="checkbox"/> Whole Blood																													
<input type="checkbox"/> Other																													