

Diagnostic reagent for quantitative in vitro determination of cholesterol in serum or plasma on Sysmex BX-Series

Order Information

Cat. No.	Kit size	Number of tests	
1 1300 99 10 970	R1 4 x 31.7 mL	BX-3010 4 x 200 tests	
		BX-4000 4 x 154 tests	

Method

"CHOD-PAP": enzymatic photometric test

Principle

Determination of cholesterol after enzymatic hydrolysis and oxidation. The colorimetric indicator is quinoneimine which is generated from 4-aminoantipyrine and phenol by hydrogen peroxide under the catalytic action of peroxidase (Trinder's reaction) [1,2].

Cholesterol ester + H_2O Cholesterol + Fatty acid

Cholesterol + O_2 <u>CHO</u> Cholesterol-3-one + H_2O_2

 $2 H_2O_2 + 4$ -Aminoantipyrine + Phenol POD Quinoneimine + $4 H_2O$

Reagent

Components and Concentrations

Good's buffer	pH 6.7	50 mmol/L
Phenol		5 mmol/L
4-Aminoantipyrine		0.3 mmol/L
Cholesterol esterase	(CHE)	\geq 200 U/L
Cholesterol oxidase	(CHO)	\geq 50 U/L
Peroxidase	(POD)	\ge 3 kU/L

Storage Instructions and Reagent Stability

The reagent is stable up to the end of the indicated month of expiry, if stored at $2 - 8^{\circ}$ C, protected from light and contamination is avoided. Do not freeze the reagent!

Warnings and Precautions

- 1. The reagent contains sodium azide (0.95 g/L) as preservative. Do not swallow! Avoid contact with skin and mucous membranes.
- N-acetylcysteine (NAC), acetaminophen and metamizole medication leads to falsely low results in patient samples.
- 3. In very rare cases, samples of patients with gammopathy might give falsified results [8].
- Please refer to the safety data sheets and take the necessary precautions for the use of laboratory reagents. For diagnostic purposes, the results should always be assessed with the patient's medical history, clinical examinations and other findings.
 For professional use only!

Waste Management

Please refer to local legal requirements.

Reagent Preparation

The reagent is ready to use. The bottles are placed directly into the reagent tray.

Specimen

Serum, heparin plasma or EDTA plasma

Stability [3]:		-	
7 days	at	20 – 25°C	
7 days	at	4 – 8°C	
3 months	at	–20°C	

Freeze only once. Discard contaminated specimens.

Calibrators and Controls

For calibration the DiaSys TruCal U calibrator is recommended. The assigned values of the calibrator have been made traceable to the reference method gas chromatography-isotope dilution mass spectrometry (GC-IDMS). For internal quality control DiaSys TruLab N and P or TruLab L controls should be assayed. Each laboratory should establish corrective actions in case of deviations in control recovery.

	Cat. No.	ŀ	Kit s	ize
TruCal U	5 9100 99 10 063	20	х	3 mL
	5 9100 99 10 064	6	х	3 mL
TruLab N	5 9000 99 10 062	20	х	5 mL
	5 9000 99 10 061	6	х	5 mL
TruLab P	5 9050 99 10 062	20	х	5 mL
	5 9050 99 10 061	6	х	5 mL
TruLab L Level 1	5 9020 99 10 065	3	х	3 mL
TruLab L Level 2	5 9030 99 10 065	3	х	3 mL

Performance Characteristics

Measuring range up to 750 mg/dL (19.4 mmol/L) cholesterol				
(in case of higher concentrations re-measure samples after manual dilution				
with NaCI (9 g/L) or use rerun function)				
Limit of detection**	1 mg/dL (0.026 mmol/L) cholesterol			
On-board stability	6 weeks			
Calibration stability	6 weeks			

Interfering substance	Interferences < 10%	Analyte concentration	
Ascorbate	up to 6 mg/dL	166 mg/dL (4.29 mmol/L)	
Hemoglobin	up to 300 mg/dL	228 mg/dL (5.90 mmol/L)	
Bilirubin, conjugated	up to 13 mg/dL	193 mg/dL (4.99 mmol/L)	
Bilirubin, unconjugated	up to 15 mg/dL	209 mg/dL (5.41 mmol/L)	
Lipemia (triglycerides)	up to 2000 mg/dL	180 mg/dL (4.66 mmol/L)	
For further information on interfering substances refer to Young DS [7].			

Precision (BX-3010)			
Within run (n=20)	Sample 1	Sample 2	Sample 3
Mean [mg/dL]	82.2	137	286
Mean [mmol/L]	2.13	3.54	7.39
Coefficient of variation [%]	2.02	1.82	1.70
Between run (n=20)	Sample 1	Sample 2	Sample 3
Mean [mg/dL]	142	225	340
Mean [mmol/L]	3.67	5.83	8.79
Coefficient of variation [%]	3.01	2.07	1.49

Method comparison (n=106)	
Test x	Cholesterol FS (BioMajesty 6010C)
Test y	Cholesterol FS (BX-3010)
Slope	1.002
Intercept	-4.04 mg/dL (-0.104 mmol/L)
Coefficient of correlation	0.999

** lowest measurable concentration which can be distinguished from zero mean + 3 SD (n=20) of an analyte free specimen

Conversion factor

Cholesterol [mg/dL] x 0.02586 = Cholesterol [mmol/L]

Reference Range [4]

Desirable	< 200 mg/dL (< 5.2 mmol/L)
Borderline high risk	200 – 240 mg/dL (5.2 – 6.2 mmol/L)
High risk	≥ 240 mg/dL (≥ 6.2 mmol/L)

Each laboratory should check if the reference ranges are transferable to its own patient population and determine own reference ranges if necessary.

Clinical Interpretation

The European Task Force on Coronary Prevention recommends to lower TC concentration to less than 190 mg/dL (5.0 mmol/L) and LDL-cholesterol to less than 115 mg/dL (3.0 mmol/L) [5].

Literature

- Artiss JD, Zak B. Measurement of cholesterol concentration. In: Rifai N, Warnick GR, Dominiczak MH, eds. Handbook of lipoprotein testing. Washington: AACC Press, 1997: p. 99-114.
- Deeg R, Ziegenhorn J. Kinetic enzymatic method for automated determination of total cholesterol in serum. Clin Chem 1983; 29: 1798-802.
- Guder WG, Zawta B et al. The Quality of Diagnostic Samples. 1st ed. Darmstadt: GIT Verlag; 2001. p. 22-3.
- Schaefer EJ, McNamara J. Overview of the diagnosis and treatment of lipid disorders. In: Rifai N, Warnick GR, Dominiczak MH, eds. Handbook of lipoprotein testing. Washington: AACC press, 1997: p. 25-48.
- Recommendation of the Second Joint Task Force of European and other Societies on Coronary Prevention. Prevention of coronary heart disease in clinical practice. Eur Heart J 1998; 19: 1434-503.
- Rifai N, Bachorik PS, Albers JJ. Lipids, lipoproteins and apolipoproteins. In: Burtis CA, Ashwood ER, editors. Tietz Textbook of Clinical Chemistry. 3rd ed. Philadelphia: W.B Saunders Company; 1999. p. 809-61.
- Young DS. Effects of Drugs on Clinical Laboratory Tests. 5th ed. Volume 1 and 2. Washington, DC: The American Association for Clinical Chemistry Press 2000.
- Bakker AJ, Mücke M. Gammopathy interference in clinical chemistry assays: mechanisms, detection and prevention. ClinChemLabMed 2007;45(9):1240-1243.

Manufacturer

CE

IVD

DiaSys Diagnostic Systems GmbH Alte Strasse 9 65558 Holzheim Germany

Chemistry Parameters 1	Sysmex BX-3010 Chemistry Analyzer
Method No * Method Name CHOL	Reagent Name Reagent (ul.) Water (ul.)
Print Name Cholesterol MethodColor	
Sample Type Serum	R2 Disable
Unit mg/dL	Diluent Disable
Assay Type End	Sample Ppt. Wash Disable
Measuring points Start End	Stirring Speed R1 Middle R2
1 <u>45</u> – <u>46</u> 2 <u>Disable</u> –	
	Normal Range No. Normal Range Name Min Max
Wave Length Prim. 510 Sec. 700	1 Male-G1 * * 2 Male-G2 * *
	3 Male-G3 * *
Newsel Constantion (cd.) Diluted Constant (cd.) Diluted	
Low Normal High	(Conc) 1 – 750
Diluent 0.0 < 1.5 < 0.0 Rerun (High/Prozone)	(mAbs/10) * *
$\Box \text{ Diluent } 0.0 < 1.5 < 0.0$	Previous Result Comparison (%) * * // *
□ Diluent 0.0 < 1.5 < 0.0	Abnormal Range (Conc) * – *
	Panic Range (Conc) * *
	Decimal Point 0 Profile SI Disable
*Entered by user	
Chemistry Parameters 2	Sysmex BX-3010 Chemistry Analyzer Analytical Parameters
Method No. * Method Name CHOL	Sample Serum
Limit Checks	Blank measurement
✓ Duplicate Limit 50 mAbs/10	Blank measurement:
✓ Sensitivity Limit 500 mAbs/10	Disable reagent blank and C1 blank
✓ Linearity Limit %	Measurement of Reagent Blank during Run: None
(mAbs/10)/min	Reagent blank measurement at calibration:
Prozone Limit Higher %	Reagent blank (No sample)
	Duplicate
SL1-S SL1-F	Reagent blank limit checks:
SL2-S – SL2-F	
Sensitivity mAbs/10	Instrument Factor
✓ Absorbance Limit Abs_in reaction Increase	a 1.00 b 0.00
Limit 25000 mAbs/10	

Chemistry Code 100 21

Calibration Registration	Sysmex BX-3010 Chemistry Analyzer Analytical Parameters
Method No. * Method Name CHOL Sample Type Serum Replication Duplicate Check Interval 42 Test without calibration Disable Calibration Type Linear	Reagent Lot No. (R1) * Last
Reagent Lot New Add	The calibration curve is lot dependent
Conc. WORK MASTER Calibr. Lot No. C1 0 Automatic entry Automatic entry C2 * Automatic entry * C3 *	II Reagent blank mAbs/10 Last Blank Automatic entry mAbs/10 Last Calibration Curve Conc Absorbance mAbs/10 Recalculation
*Entered by user	

Chemistry Code 100 21

Chemistry Parameters	Sysm	ex BX-4000 Cho Analy	emistry Analyzer tical Parameters
Method * Name CHOL	Reagent Name	Reagent (µL)	Water (µL)
Print Name Cholesterol R1	CHOL	180	
Sample Serum R2 ✓	Enable		
Unit mg/dL			
Assay Type End Diluent I] Enable		
Measuring points Start End Decimal	Points 0		
1 67 – 68			
🗆 Enable 2 📃 –			
Norm No.	l Range Normal Range Name	Min	Max
Wave Length 1 Prim. 510 Sec □ Disable 700 2	Male-G1 Male-G2	*	*
3	Male-G3 Female-G1	*	*
Normal Sampling Sample (uL) Diluent (uL)	Technical Range		
Dilution 2.0	(Conc)	1	750
Dilution			lI
Dilution		Reagent Name	
	SPT Wash		
	Stirring Speed R1	Middle R2	
*Entered by user			
Entered by user			
Chemistry Parameters	Sysm	ex BX-4000 Che Analy	emistry Analyzer rtical Parameters
Method No. * Name CHOL Sample Serum]		
Limit Checks	Blank measurement		
Duplicate Limit 50 mAbs/10	Blank measurement: Disable reagent blank and S1 k	olank	
✓ Sensitivity Limit 500 mAbs/10	Measurement of Reagent Blank	during Run:	
✓ Linearity Limit % (mAbs/10)/min	None		
Prozone Limit % Upper	Reagent blank measurement at o	calibration:	
SL1-S SL1-F	The number of measurement:		J
SL2-S SL2-F	Duplicate		
Sensitivity mAbs/10	Reagent blank limit checks:	20	
✓ Absorbance Limit		20	
Reaction	Instrument Factor		
Limit 25000 mAbs/10	a 1.00	b 0.00	
	II		

Chemistry Code 100 21

Registration Calibration	Sysmex BX-4000 Chemistry Analyzer Analytical Parameters
Method * Name CHOL	R Lot No. R1 * Last
Sample Serum	
Sampling Duplicate	Master w015 Work
Check Interval 42 days	
Auto Change Lot Full Calibration	1956
Auto Interval hours	1384
Type Linear Lot New	280
Material Name NaCL/TruCal U	The calibration curve is lot dependent
	Reagent blank mAbs/10 Last
S1 0 Automatic entry Automatic entry	Blank Automatic entry mAbs/10 Last
S2 *	Type Conc.
S4	Absorbance mAbs/10 Recalculation
S6 ^	
K Automatic entry S1 Blank Reagent Blank for S1	
*Entered by user	